

Nonparametric Inference with SnPM

Thomas Nichols, Ph.D.
Department of Statistics
Warwick Manufacturing Group
University of Warwick

Vancouver SPM Course
August 7, 2010

Outline


- Orientation
- Assessing Statistic images
- The Multiple Testing Problem
- Random Field Theory & FWE
- Permutation & FWE
- False Discovery Rate

Outline


- Orientation
- Assessing Statistic images
- The Multiple Testing Problem
- Random Field Theory & FWE
- Permutation & FWE
- False Discovery Rate

Nonparametric Inference

- Parametric methods
 - Assume distribution of statistic under null hypothesis
 - Needed to find P-values, u_α
- Nonparametric methods
 - Use *data* to find distribution of statistic under null hypothesis
 - Any statistic!



Parametric Null Distribution



Nonparametric Null Distribution

Permutation Test Toy Example

- Data from V1 voxel in visual stim. experiment
 - A: Active, flashing checkerboard B: Baseline, fixation
 - 6 blocks, ABABAB Just consider block averages...

A	B	A	B	A	B
103.00	90.48	99.93	87.83	99.76	96.06

- Null hypothesis H_0
 - No experimental effect, A & B labels arbitrary
- Statistic
 - Mean difference

Permutation Test Toy Example

- Under H_0
 - Consider all equivalent relabelings

AAABBB	ABABAB	BAAABB	BABBAA
AABABB	ABABBA	BAABAB	BBAAAB
AABBAB	ABBAAB	BAABBA	BBAABA
AABBBA	ABBABA	BABAAB	BBBABAA
ABAABB	ABBBAA	BABABA	BBBAAA

Permutation Test Toy Example

- Under H_0
 - Consider all equivalent relabelings
 - Compute all possible statistic values

AAABBB	4.82	ABABAB	9.45	BAAABB	-1.48	BABBAA	-6.86
AABABB	-3.25	ABABBA	6.97	BAABAB	1.10	BBAAAB	3.15
AABBAB	-0.67	ABBAAB	1.38	BAABBA	-1.38	BBAABA	0.67
AABBBA	-3.15	ABBABA	-1.10	BABAAB	-6.97	BBABAA	3.25
ABAABB	6.86	ABBBAA	1.48	BABABA	-9.45	BBBAAA	-4.82

Permutation Test Toy Example

- Under H_o
 - Consider all equivalent relabelings
 - Compute all possible statistic values
 - Find 95%ile of permutation distribution

AAABBB	4.82	ABABAB	9.45	BAAABB	-1.48	BABBAA	-6.86
AABABB	-3.25	ABABBA	6.97	BAABAB	1.10	BBAAAB	3.15
AABBAB	-0.67	ABBAAB	1.38	BAABBA	-1.38	BBAABA	0.67
AABBBA	-3.15	ABBABA	-1.10	BABAAB	-6.97	BBABAA	3.25
ABAABB	6.86	ABBBAA	1.48	BABABA	-9.45	BBBAAA	-4.82


Permutation Test Toy Example

- Under H_o
 - Consider all equivalent relabelings
 - Compute all possible statistic values
 - Find 95%ile of permutation distribution

AAABBB	4.82	ABABAB	9.45	BAAABB	-1.48	BABBAA	-6.86
AABABB	-3.25	ABABBA	6.97	BAABAB	1.10	BBAAAB	3.15
AABBAB	-0.67	ABBAAB	1.38	BAABBA	-1.38	BBAABA	0.67
AABBBA	-3.15	ABBABA	-1.10	BABAAB	-6.97	BBABAA	3.25
ABAABB	6.86	ABBBAA	1.48	BABABA	-9.45	BBBAAA	-4.82



Permutation Test Toy Example

- Under H_0
 - Consider all equivalent relabelings
 - Compute all possible statistic values
 - Find 95%ile of permutation distribution



Controlling FWE: Permutation Test


- Parametric methods
 - Assume distribution of *max* statistic under null hypothesis
- Nonparametric methods
 - Use *data* to find distribution of *max* statistic under null hypothesis
 - Again, any *max* statistic!



Permutation Test

Smoothed Variance t


- Collect max distribution
 - To find threshold that controls FWE
- Consider smoothed variance t statistic



Permutation Test

Smoothed Variance t

- Collect max distribution
 - To find threshold that controls FWE
- Consider smoothed variance t statistic



Permutation Test Strengths



- Requires only assumption of exchangeability
 - Under H_0 , distribution unperturbed by permutation
 - Allows us to build permutation distribution
- Subjects are exchangeable
 - Under H_0 , each subject's A/B labels can be flipped
- fMRI scans not exchangeable under H_0
 - Due to temporal autocorrelation

Permutation Test Limitations

- Computational Intensity
 - Analysis repeated for each relabeling
 - Not so bad on modern hardware
 - No analysis discussed below took more than 3 hours
- Implementation Generality
 - Each experimental design type needs unique code to generate permutations
 - Not so bad for population inference with t-tests


Permutation Test Example

- fMRI Study of Working Memory
 - 12 subjects, block design Marshuetz et al (2000)
 - Item Recognition
 - Active: View five letters, 2s pause, view probe letter, respond
 - Baseline: View XXXXX, 2s pause, view Y or N, respond
- Second Level RFX
 - Difference image, A-B constructed for each subject
 - One sample, smoothed variance t test




Permutation Test Example

- Permute!
 - $2^{12} = 4,096$ ways to flip 12 A/B labels
 - For each, note maximum of t image




Permutation Distribution
Maximum t




Maximum Intensity Projection
Thresholded t

Permutation Test Example


- Compare with Bonferroni
 - $\alpha = 0.05/110,776$
- Compare with parametric RFT
 - 110,776 2x2x2mm voxels
 - 5.1x5.8x6.9mm FWHM smoothness
 - 462.9 RESELs




t_{11} Statistic, Nonparametric Threshold



t_{11} Statistic, RF & Bonf. Threshold



Test Level vs. t_{11} Threshold



Smoothed Variance t Statistic,
Nonparametric Threshold ¹⁹

Does this Generalize? RFT vs Bonf. vs Perm.

	df	<i>t</i> Threshold (0.05 Corrected)		
		RF	Bonf	Perm
Verbal Fluency	4	4701.32	42.59	10.14
Location Switching	9	11.17	9.07	5.83
Task Switching	9	10.79	10.35	5.10
Faces: Main Effect	11	10.43	9.07	7.92
Faces: Interaction	11	10.70	9.07	8.26
Item Recognition	11	9.87	9.80	7.67
Visual Motion	11	11.07	8.92	8.40
Emotional Pictures	12	8.48	8.41	7.15
Pain: Warning	22	5.93	6.05	4.99
Pain: Anticipation	22	5.87	6.05	5.05



RFT vs Bonf. vs Perm.

	df	No. Significant Voxels (0.05 Corrected)			SmVar t Perm
		t	Bonf	Perm	
Verbal Fluency	4	0	0	0	0
Location Switching	9	0	0	158	354
Task Switching	9	4	6	2241	3447
Faces: Main Effect	11	127	371	917	4088
Faces: Interaction	11	0	0	0	0
Item Recognition	11	5	5	58	378
Visual Motion	11	626	1260	1480	4064
Emotional Pictures	12	0	0	0	7
Pain: Warning	22	127	116	221	347
Pain: Anticipation	22	74	55	182	402



Using SnPM to Assess Reliability with Small Groups

- Consider n=50 group study
 - Event-related Odd-Ball paradigm, Kiehl, et al.
- Analyze all 50
 - Analyze with SPM and SnPM, find FWE thresh.
- Randomly partition into 5 groups 10
 - Analyze each with SPM & SnPM, find FWE thresh
- Compare reliability of small groups with full
 - With and without variance smoothing



SPM t_{11} : 5 groups of 10 vs all 50 5% FWE Threshold



SnPM t : 5 groups of 10 vs. all 50 5% FWE Threshold



SnPM SmVar t : 5 groups of 10 vs. all 50 5% FWE Threshold



Conclusions


- Nonparametric Permutation
 - Good when Normality is question
 - Good with tiny group inference & variance smoothing
- Come to practical for more!

Using SnPM

- Choose right ‘Plug in’
 - Depends on your design
- Cluster size inference?
 - Yes: Commit to cluster-forming threshold now
 - Yes: Don’t commit, collect *huge* SnPM_ST file
- Approximate Test
 - Are you in a hurry?
 - Or are more than 10,000 perms possible?
 - Then ‘Yes’ approximate, and choose 1k-10k perms
- Which image? T or P?
 - T: Traditional voxel-wise or cluster-wise FWE
 - P: Uncorrected voxel-wise or FDR


Henson et al. Faces Data

- Famous-vs-Nonfamous faces
 - Chapter 30 of SPM manual
 - Main effect, Any Faces – Checkerboard
 - 12 subjects
 - ‘cons_can’ Canonical HRF only
 - ‘cons_informed’ Canonical + Temp Deriv + Disp Deriv
- Will compare SnPM to SPM
 - For 1-sample t-test (cons_can) & Repeated measures ANOVA (cons_informed)




Voxel-Wise Results Canonical HRF t test


- SPM
 - $u_{FWE} = 9.071$, 371 voxels



- SnPM
 - $u_{FWE} = 7.925$, 917 voxels



- SnPM w/ Var Smoothing
 - (u_{FWE} not comparable) 3575 voxels w/ 6mm
 - 3483 voxels w/ 4mm



Voxel-Wise Results

3-Basis F test

- SPM
 - u_{FWE} : 3733 voxels
- SnPM
 - u_{FWE} : 5105 voxels

